3.5625”

4.9375”

5.1875”

ORTHO®

FLOWER, FRUIT & VEGETABLE

INSECT KILLER

READY-SPRAY® II *INSECTICIDA PARA FLORES, FRUTAS Y VEGETALES*

KILLS 100+ *LISTED* **INSECTS FAST**

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel booklet for additional precautionary statements.

MANTENGA FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCION Consulte el folleto del panel posterior para conocer los avisos de precaución adicionales.

FOR RESIDENTIAL OUTDOOR USE SITES ONLY
NOT FOR COMMERCIAL PRODUCTION

Active Ingredient	
Acetamiprid	0.5%
Other Ingredients	99.5%
Total	100.0%

FPO

LB2622

NET CONTENTS/CONTENIDO NETO
32 fl oz (946 ml)

3.8125”

EPA Reg No. 8033-107-239 (32 oz)
Ortho Flower, Fruit & Vegetable Insect Killer Ready-Spray II

2.6875”

2.875”

4.8125”

5.1875”

RIESGOS AMBIENTALES

Para proteger el medio ambiente, no permita que el pesticida ingrese o discurra al desagües de tormenta, zanjas de drenaje, canalones o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no haya un pronóstico de lluvia para las próximas 24 horas, asegurará que el viento o la lluvia no soplen o laven el pesticida del área de tratamiento. Enjuagar el equipo de aplicación en la zona tratada evitará que el producto discurra a cuerpos de agua o sistemas de drenaje. Este pesticida es tóxico para animales y plantas silvestres. Este producto es tóxico para las abejas expuestas al tratamiento de manera directa. No aplique este producto mientras las abejas visitan activamente el área tratada.

RIESGOS FISICOS O QUIMICOS: No aplique este producto sobre equipos eléctricos porque puede recibir una descarga eléctrica.

AVISO: En la medida que la ley lo permita, el comprador asume todos los riesgos de uso, almacenamiento o manejo de este producto si no se siguen las instrucciones correspondientes.

†**La Garantía ORTHO:** Si por algún motivo, usted, el consumidor, no está satisfecho con este producto, envíenos el comprobante original de compra por correo para recibir el reembolso total del precio de compra.

24

ORTHO®

FLOWER, FRUIT & VEGETABLE

INSECT KILLER

READY-SPRAY® II

OPEN Resealable Label for Directions & Precautions
Español ver página 12

WON’T HARM PLANTS OR BLOOMS!
Simply Attach to Hose and Go

KILLS 100+ LISTED INSECTS FAST

Aphids · Caterpillars · Thrips · Whiteflies · Japanese Beetles

WHERE TO USE:

Roses & Flowers · Shrubs · Listed Fruits & Vegetables

WHEN TO APPLY:

Apply as soon as insect problems are noticed.

F.P.O.
80%

0 71549 03460 3
1005

Questions, Comments or Medical Information
Preguntas, comentarios o información médica
1-800-225-2883

©2012 The ORTHO Group. World rights reserved.
Sold by **The ORTHO Group**
P.O. Box 190, Marysville, OH 43040
Product of Japan.
EPA Reg. No. 8033-107-239
EPA Est. 239-IA-3[I], 239-MS-001[M], 58996-MO-1[A], 085652-OH-1[W]
Superscript is first letter of lot number.

FPO

LB2623

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT APPLICATION INFORMATION

- For outdoor residential use sites only.
- Not for commercial production.
- Do not apply this product in a way that the spray will contact persons, either directly or through drift.
- Keep children and pets out of treated areas until sprays have thoroughly dried.
- When applying outdoors:
  - Do not apply if wind speed exceeds 10 mph.
  - Always stand “up wind” when applying this product.

READY-SPRAY® II HOSE END APPLICATOR

Shake container gently before using.

Connect:	• Connect sprayer to hose. • Turn on water.
Spray: 	• To begin spraying, point nozzle in the direction you want to spray making sure that the end of the nozzle is 8 to 12 inches away from the plants. • Slide thumb activation switch forward to **ON** position. • Direct spray toward the upper and lower leaf surfaces and stems where pests appear. • Spray until the surface of the foliage is wet.
Finish: 	• To **STOP** spraying, slide thumb activation switch to **OFF** position. • Turn off water. • Relieve water pressure by sliding thumb switch to **WATER** position until water slows to a drip. Then, slide switch back to **OFF** position. • Disconnect the sprayer from hose. **Do not use hose end applicator indoors.**

GARDENS

TO KILL THESE INSECTS	Aphids (including Green Peach, Wooly, Melon, and Cotton), Army Worms (including Fall Army Worm, and Southern Army Worm), Banded Grape Bug, Beetles (including Bean Leaf Beetle, Colorado Potato Beetle, Flea Beetle, Japanese Beetle, Mexican Bean Beetle and Oriental Beetle), Cucumber Beetle (including Spotted, Striped and Western Striped), Bollworm, Borers (including Dogwood Borer, European Corn Borer, Peach Twig Borer and Squash Vine Borer), Budworm (including Tobacco Budworm), Caterpillars (including Red Humped Caterpillar, Tomato Fruitworm, and Cabbage Looper), Rose Chafer, Plum Curculio, European Chafer, European Pine Sawfly, Filbertworm, Flea Hopper, Fruit Fly (including Cherry, Black Cherry and Western Cherry), Western Grapeleaf Skeletonizer, Hickory Shuckworm, Hornworms, Katydid, Phylloxera (aerial form only), Leafhoppers (including Glassy Wing Sharpshooter and Potato Leafhopper), Leaf Miners (including Chrysanthemum, Citrus, and Tentiform), Lesser Apple Worm, Apple Maggot, Mealybugs (including Citrus, Grape, Obscure, Longtail, Pink Hybiscus, and Maderia), Melonworm, Moths (including Codling, Diamondback Moth, Gypsy Moths and Oriental Fruit Moth), Navel Orangeworm, Pecan Nut Casebearer, Pickleworm, Plantbugs (including Tarnished, and Mullein Plantbug), Psyllids, European Apple Sawfly, Scale (including Asian Cycad Scale, Caribbean Black Scale, California Red Scale, Citricola Scale, Cottony Maple Scale, Euonymus Scale, Fletcher Scale, Florida Wax Scale, Green Shield Scale, Indian Wax Scale, Oyster Shell Scale, Pine Needle Scale, Red Scale, San Jose Scale and Tea Scale), Squash Bug, Stinkbug, Swede Midge, Thrips (including Citrus, Cotton, Palm, and Western Flower), Weevils (including Pecan Weevil, Pepper Weevil and Strawberry Weevil), Whiteflies (including Greenhouse, Sweet Potato, Silverleaf, Banded and Giant)
ON THESE PLANTS	Ornamentals, Roses, Flowers, Trees, Shrubs
WHEN TO APPLY	Apply as soon as insect problems are noticed. Wait 1 to 4 weeks to re-apply if insect problems reappear. Do not apply more than 5 times in a season.

3.125”

3.125”

EPA Reg No. 8033-107-239 (32 oz)
Ortho Flower, Fruit & Vegetable Insect Killer Ready-Spray II

2.6875”

2.6875”

4.8125”

5.1875”

EDIBLES	
TO KILL THESE INSECTS	Aphids (including Green Peach, Wooly, Melon, and Cotton), Army Worms (including Fall Army Worm, and Southern Army Worm), Asian Citrus Psyllid, Banded Grape Bug, Beetles (including Bean Leaf Beetle, Colorado Potato Beetle, Flea Beetle, Japanese Beetle, Mexican Bean Beetle, Sap Beetles, and Oriental Beetle), Blueberry Maggot, Blueberry Gall Midge, Cucumber Beetle (including Spotted, Striped and Western Striped), Budworm (including Tobacco Budworm), Caterpillars (including Red Humped Caterpillar, Tomato Fruitworm, Cherry Fruitworm, Cranberry Fruitworm, Cranberry Tipworm, Sparganothis Fruitworm, Western Raspberry Fruitworm [adult] and Cabbage Looper), Rose Chafer, Plum Curculio, Filbertworm, Flea Hopper, Fruit Fly (including Cherry, Black Cherry and Western Cherry), Hickory Shuck Worm, Western Grapeleaf Skeletonizer, Hornworms, Katydid, Phylloxera (aerial form only), Grape Cane Girdler, Leafhoppers (including Glassy Wing Sharpshooter, Grape, Variegated, and Potato Leafhopper), Leaf Miners (including Chrysanthemum, Citrus, and Tentiform), Lesser Apple Worm, Apple Maggot, Mealybugs (including Citrus, Gill’s, Grape, Obscure, Longtail, Pink Hybiscus, and Maderia), Melonworm, Moths (including Codling, Diamondback Moth, Gypsy Moths and Oriental Fruit Moth), Oblique Banded Leaf Roller, Navel Orangeworm, Peach Twig Borer, Pecan Nut Casebearer, Pickleworm, Plantbugs (including Tarnished, and Mullein Plantbug), Psyllids, European Apple Sawfly, Scale (including Asian Cycad Scale, Caribbean Black Scale, California Red Scale, Citricola Scale, Cottony Maple Scale, Euonymus Scale, Fletcher Scale, Florida Wax Scale, Green Shield Scale, Indian Wax Scale, Oyster Shell Scale, Pine Needle Scale, Red Scale, San Jose Scale and Tea Scale), Spanworm, Spittlebug, Squash Bug, Squash Vine Borer, Stinkbug, Strawberry Rootworm, Thrips (including Citrus, Cotton, Palm, and Western Flower), Walnut Husk Fly, Weevils (including Pecan Weevil, Pepper Weevil and Strawberry Weevil), Whiteflies (including Greenhouse, Sweet Potato, Silverleaf, Banded and Giant)

4

EDIBLES	
ON THESE PLANTS	**Vegetables:** Amaranth, Arrugula, Broccoli, Brussels Sprouts, Cabbage, Cardoon, Cauliflower, Celery, Celtuce, Chervill, Chrysanthemum (edible leaved, garland), Collards, Corn Salad, Cress (garden, upland), Dandelion, Dock, Eggplant, Endive, Florence Fennel, Groundcherry, Kale, Kohlrabi, Lettuce (head, leaf), Mizuna, Mustard Greens, Mustard Spinach, Orach, Parsley, Pepino, Pepper (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Purslane (garden, winter), Radicchio, Rape Greens, Rhubarb, Spinach (leaf, vine, New Zealand), Swiss Chard, Tomatillo, Tomato **Citrus Fruits:** Calamondin, Citrus hybrids (chironja, tangelo, tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sour, sweet), Pummelo, Satsuma mandarin **Pome Fruits:** Apple, Crabapple, Loquat, Mayhaw, Pear, Quince **Grapes and Other Climbing Vine Small Fruit (except fuzzy kiwifruit):** Gooseberry, Hardy Kiwifruit, Amur River Grape, Maypop, Schisandra Berry **Tuberous and Corm Vegetables:** Potato, Sweet Potato, Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible Canna, Cassava (Bitter and Sweet), Chayote (Root), Chufa, Sasheen, Ginger, Leren, Tanier, Tumeric, Yam Bean, True Yam **Stone Fruit:** Apricot, Cherry (Sweet and Tart), Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot, Prune (Fresh) **Cucurbits:** Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible), *Mormordica* spp., Muskmelon (hybrid and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Squash (summer and winter), Watermelon **Tree Nuts:** Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia (bush nut), Pecan, Pistachio, Walnut (black and English [Persian])

5

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

EDIBLES	
ON THESE PLANTS (cont’d.)	**Edible Podded Legume Vegetables and Succulent Shelled Peas and Beans:** Bean (*Phaseolus* spp.) (includes runner bean, snap bean, wax bean, lima bean [green]); Bean (*Vigna* spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean, blackeyed pea, cowpea, southern pea); Jackbean; Broad Bean (succulent); Pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, snow pea, sugar snap pea, English pea, garden pea, green pea); Soybean (immature seed); Sword Bean; and Pigeon Pea **Strawberries and Other Low Growing Berries:** Bearberry; Bilberry; Lowbush Blueberry; Cloudberry; Cranberry; Lingonberry; Muntries; Partridgeberry; and cultivars, varieties, and/or hybrids of these **Blueberries and Other Bush and Cane Berries:** Aronia berry; blueberry, highbush and lowbush; buffalo currant; Chilean guava; currant, red and black; elderberry; European barberry; gooseberry; cranberry, highbush; edible honeysuckle; huckleberry; jostaberry; Juneberry; lingonberry; natie currant; salal; sea buckthorn; and cultivars, varieties, and/or hybrids of these; and, Blackberry, Loganberry, Raspberry (black and red), wild raspberry, and cultivars, varieties, and/or hybrids of these **Onions and Other Bulb Vegetables:** Chives, fresh leaves; Chinese chives, fresh leaves; daylily bulbs, Elegans hosta, Fritillaria leaves and bulbs; bulb garlic; great headed bulb garlic, serpent bulb garlic; kurrat; lady’s leek; leek; wild leek; lily bulb; Beltsville bunching onion; bulb onion; Chinese bulb onion; fresh onion; green onion; macrostem onion; pearl onion; potato bulb onion; treetops onion; Welsh onion tops; shallot bulb and fresh leaves; and cultivars, varieties, and/or hybrids of these
WHEN TO APPLY	Apply as soon as insect problems are noticed.

6

	Days to Wait to Harvest	Days to Wait Before Reapplication	Maximum Number of Applications per Season
Amaranth, Arrugula, Broccoli, Brussels Sprouts, Cabbage, Cardoon, Cauliflower, Celery, Celtuce, Chervill, Chrysanthemum (edible leaved, garland), Collards, Corn Salad, Cress (garden, upland), Dandelion, Dock, Eggplant, Endive, Florence Fennel, Groundcherry, Kale, Kohlrabi, Lettuce (head, leaf), Mizuna, Mustard Greens, Mustard Spinach, Orach, Parsley, Pepino, Pepper (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Purslane (garden, winter), Radicchio, Rape Greens, Rhubarb, Spinach (leaf, vine, New Zealand), Swiss Chard, Tomatillo, Tomato, Potato, Sweet Potato, Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible Canna, Cassava (Bitter and Sweet), Chayote (Root), Chufa, Sasheen, Ginger, Leren, Tanier, Tumeric, Yam Bean, True Yam, Chives, fresh leaves; Chinese chives, fresh leaves; daylily bulbs, Elegans hosta, Fritillaria leaves and bulbs; bulb garlic; great headed bulb garlic, serpent bulb garlic; kurrat; lady’s leek; leek; wild leek; lily bulb; Beltsville bunching onion; bulb onion; Chinese bulb onion; fresh onion; green onion; macrostem onion; pearl onion; potato bulb onion; treetops onion; Welsh onion tops; shallot bulb and fresh leaves; and cultivars, varieties, and/or hybrids of these	7	7	4

7

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

	Days to Wait to Harvest	Days to Wait Before Reapplication	Maximum Number of Applications per Season
Citrus Fruits: Calamondin, Citrus hybrids (chironja, tangelo, tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sour, sweet), Pummelo, Satsuma mandarin	7	7	5
Pome Fruits: Apple, Crabapple, Loquat, Mayhaw, Pear, Quince	7	12	4
Grapes and Other Climbing Vine Small Fruits (except fuzzy kiwifruit): Amur river grape; gooseberry, hardy kiwifruit, Maypop, schisandra berry, and cultivars, varieties, and/or hybrids of these	7	14	2
Stone Fruit: Apricot, Cherry (Sweet and Tart), Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot, Prune (Fresh)	7	10	4
Cucurbits: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible), *Mormordica* spp., Muskmelon (hybrid and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Squash (summer and winter), Watermelon	0	5	5

8

	Days to Wait to Harvest	Days to Wait Before Reapplication	Maximum Number of Applications per Season
Tree Nuts: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia (bush nut), Pecan, Pistachio, Walnut (black and English [Persian])	14	14	4
Edible Podded Legume Vegetables and Succulent Shelled Peas and Beans: Bean (*Phaseolus* spp.) (includes runner bean, snap bean, wax bean, lima bean [green]); Bean (*Vigna* spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean, blackeyed pea, cowpea, southern pea); Jackbean; Broad Bean (succulent); Pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, snow pea, sugar snap pea, English pea, garden pea, green pea); Soybean (immature seed); Sword Bean; and Pigeon Pea	7	7	3
Strawberries and Other Low Growing Berries: Bearberry; Bilberry; Lowbush Blueberry; Cloudberry; Cranberry; Lingonberry; Muntries; Partridgeberry; and cultivars, varieties, and/or hybrids of these	1	7	2
Blueberries and Other Bush and Cane Berries: Aronia berry; blueberry, highbush and lowbush; buffalo currant; Chilean guava; currant, red and black; elderberry; European barberry; gooseberry; cranberry, highbush; edible honeysuckle; huckleberry; jostaberry; Juneberry; lingonberry; natie currant; salal; sea buckthorn; and cultivars, varieties, and/or hybrids of these; and, Blackberry, Loganberry, Raspberry (black and red), wild raspberry, and cultivars, varieties, and/or hybrids of these	1	7	5

9

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

STORAGE AND DISPOSAL

STORAGE	Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place; preferably in a locked storage area.
DISPOSAL	**Non-refillable container. Do not reuse or refill this container.** **If empty**: Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

IF ON SKIN	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	• Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have the person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

10

For Medical Emergencies, Call 24 Hours a Day **1-800-225-2883**
Have the product container or label with you when calling a poison control center or doctor or going for treatment.
You may also contact 1-800-225-2883 for emergency medical treatment information.

NOTE TO PHYSICIAN: There is no specific antidote. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to other materials other than this product may have occurred.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This pesticide is toxic to wildlife. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treated area.

PHYSICAL OR CHEMICAL HAZARDS: Do not apply this product in or on electrical equipment due to the possibility of shock hazard.

NOTICE: To the extent allowed by law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

†THE ORTHO GUARANTEE: If for any reason you, the consumer, are not satisfied with this product, mail us original proof of purchase to obtain a full refund of your purchase price.

11

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

INFORMACION DE APLICACION IMPORTANTE

- Unicamente para uso residencial en exteriores.
- No lo use para aplicaciones comerciales.
- No aplique este producto de manera que entre en contacto directo o por deriva con personas.
- Mantenga a los niños y mascotas lejos del área tratada hasta que el rociado se haya secado por completo.
- Cuando se aplica en exteriores:
  - No aplique si la velocidad del viento es mayor que 16 km/h.
  - Siempre colóquese de espaldas al viento cuando aplique este producto.

ROCIADOR DE MANGUERA READY-SPRAY® II

Agite el envase suavemente antes de usar.

Conectar:	• Conecte el rociador a la manguera. • Encienda el agua.
Rociar: 	• Para empezar a rociar, dirija la boquilla en la dirección que desea rociar, asegurándose de que la punta de la boquilla esté de 20 cm a 30 cm de las plantas. • Deslice con el pulgar la llave de activación hacia delante, a la posición de **ENCENDIDO (ON)**. • Dirija el rocío hacia la superficie superior e inferior de las hojas y los tallos donde aparecen las plagas. • Rocíe solamente hasta que la superficie de la planta esté húmeda.
Finalizar: 	• Para **DETENER** el rociado, deslice con el pulgar la llave de activación a la posición de **APAGADO (OFF)**. • Apague el agua. • Disminuya la presión del agua deslizando la llave con el pulgar hacia la posición de **AGUA (WATER)** hasta que el agua se reduzca a un goteo. Luego, deslice la llave de regreso hacia la posición **APAGADO (OFF)**. • Desconecte el rociador de la manguera. **No utilice el rociador de manguera en interiores.**

JARDINES

PARA ELIMINAR ESTOS INSECTOS	Áfidos (incluidos el verde del durazno, el lanoso, el del melón y el algodonoso), gusanos del ejército (incluidos del ejército del otoño y del ejército del sur), insectos de la vid con bandas, escarabajos (incluidos el escarabajo de la hoja del frijol, el escarabajo de la patata de Colorado, el escarabajo pulga, el escarabajo japonés, el escarabajo mexicano del frijol y el escarabajo oriental), escarabajos del pepino (incluidos el manchado, el rayado y el rayado del oeste), larvas de polilla, barrenadores (incluidos del cornejo, del maíz europeo, del peral y de la calabaza), cogolleros (incluido el cogollero del tabaco), orugas (incluidas la oruga roja jorobad, el gusano del tomate y el gusano falso medidor), gusarapos de la rosa, curculios de la ciruela, gusarapo europeo, tentredinidos del pino europeo, gusanos de la avellana, pulga saltarina, mosca de la fruta (incluidas del cerezo, del cerezo negro y del cerezo del oeste), esqueletizador de la hoja de la uva del oeste, barrenadores del ruezno, gusanos de cuerno, cigarra, filoxeras (sólo la forma aérea), cicadélidos (incluidos homalodisca coagulata y cicadélido de la papa), barrenadores de hojas tentiformes (incluidas los de crisantemos, de cítricos y tentiformes), gusanos del manzano pequeño, gusanos de la manzana, cochinillas harinosas (incluidas las de cítricos, de uvas, las oscuras, las de cola larga, las rosadas y las de Madeira), gusano del melón, polillas (incluidas la carpocapsa, la polilla dorso de diamante, la polilla gitana y la polilla de la fruta oriental), gusanos naranja de ombligo, gusanos barrenador de la nuez, gusanos barrenador del fruto, gorgojos de huerta (incluidas chinches lygus y gordobolo), psílidos, mosca de sierra de la manzana europea, cochinillas (incluidas la cochinilla asiática de Cycad, cochinilla negra caribeña, cochinilla roja de California, cochinilla citrícola, cochinilla algodonosa del arce, cochinilla del evónimo, cochinilla Fletcher, cochinilla de la cera de Florida, cochinilla de caparazón verde, cochinilla de la cera de la India, cochinillas concha de ostra, cochinillas de agujas de pino, cochinillas rojas, cochinilla de San José y cochinilla Tea), chinches de la calabaza, pentatómidos, mosquito de la col, trips (incluidos de cítricos, del algodón, de la palmera y de la flor del oeste), gorgojos (incluidos gorgojo de la nuez pacana, gorgojo del pimiento y gorgojo de la fresa), moscas blancas (incluidas de invernadero, de la patata dulce, de hoja plateada, con bandas y la gigante)

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

JARDINES	
EN ESTAS PLANTAS	Rosas, flores, árboles, arbustos y plantas ornamentales
CUANDO APLICAR	Aplique tan pronto como detecte los problemas de insectos. Espere entre 1 y 4 semanas para aplicar nuevamente si vuelven a aparecer los problemas de insectos. No haga más de 5 aplicaciones por temporada.

COMESTIBLES	
PARA ELIMINAR ESTOS INSECTOS	Áfidos (incluidos el verde del durazno, el lanoso, el del melón y el algodonoso), gusanos del ejército (incluidos del ejército del otoño y del ejército del sur), sílidos asiáticos de cítricos, insectos de la vid con bandas, escarabajos (incluidos el escarabajo de la hoja del frijol, el escarabajo de la patata de Colorado, el escarabajo pulga, el escarabajo japonés, el escarabajo mexicano del frijol, el escarabajo de la savia y el escarabajo oriental), gusanos de las grosellas, moscas de las agallas de las grosellas, escarabajos del pepino (incluidos el usano del arándano, gusano de la punta del arándano [Diptera: Cecidomyiidae] gusano Sapaganothis, gusano de la frambuesa occidental [adulto] y el gusanos falso medidor), gusarapos de la rosa, curculios de la ciruela, gusanos de la avellana, pulga saltarina, mosca de la fruta (incluidas del cerezo, del cerezo negro y del cerezo del oeste), barrenadores del ruezno, esqueletizador de la hoja de la uva del oeste, gusanos de cuerno, cigarra, filoxeras (sólo la forma aérea), Ampeloglypter ampelopsis, cicadélidos (incluidos homalodisca coagulata, de la uva, variegado y cicadélido de la papa), barrenadores de hojas (incluidas los de crisantemos, de cítricos y tentiformes), gusanos del manzano pequeño, gusanos de la manzana, cochinillas harinosas (incluidas las de cítricos, de uvas, las oscuras, las de cola larga, las rosadas y las de Madeira), gusano del melón, polillas (incluidas la carpocapsa, la polilla dorso de diamante, la polilla gitana y la polilla de la fruta oriental), gusanos naranja umbilicada, barrenador del melocotón, gusanos barrenador de la nuez, gusanos barrenador del fruto, gorgojos de huerta (incluidas chinches lygus y

14

COMESTIBLES	
PARA ELIMINAR ESTOS INSECTOS (cont.)	gordobolo), psílidos, mosca de sierra de la manzana europea, cochinillas (incluidas la cochinilla asiática de Cycad, cochinilla negra caribeña, cochinilla roja de California, cochinilla citrícola, cochinilla algodonosa del arce, cochinilla del evónimo, cochinilla Fletcher, cochinilla de la cera de Florida, cochinilla de caparazón verde, cochinilla de la cera de la India, cochinillas concha de ostra, cochinillas de agujas de pino, cochinillas rojas, cochinilla de San José y cochinilla Tea), gusano falso medidor, gusanos escupidores, chinches de la calabaza, barrenadores de la calabaza, gusanos de la raíz de las fresas, pentatómidos, mosquito de la col, trips (incluidos de cítricos, del algodón, de la palmera y de la flor del oeste), moscas de la nuez, gorgojos (incluidos gorgojo de la nuez pacana, gorgojo del pimiento y gorgojo de la fresa), moscas blancas (incluidas de invernadero, de la patata dulce, de hoja plateada, con bandas y la gigante)
EN ESTAS PLANTAS	**Verduras:** Amaranto, rúcula silvestre, brócoli, col de Bruselas, repollo, cardo, coliflor, apio, celtuce, perifollo, crisantemos (de hoja comestible, guirnalda), col verde, maíz salado, draba (jardín, meseta), diente de león, romaza, berenjena, endivia, hinojo de Florencia, uchuva, coles rizadas, colirrábano, lechuga (cabeza, hoja), mizuna, hojas verdes de mostaza, mostaza espinaca, plantas de sal, perejil, pepino, pimiento (morrón, chile, pimiento de cocina, pimento, dulce), verdolaga (jardín, invierno), radicheta, rape greens (especie de col), ruibarbo, espinaca (hoja, de la vid, de Nueva Zelanda), acelga, tomatillo, tomate **Frutas cítricas:** Calamondín, híbridos de cítricos (chironja, tangelo, tangor), toronja, quinito, limón, lima, mandarina (tangerina), naranja (agria, dulce), pomelo, mandarina Satsuma **Frutas de pepitas:** Manzana, manzano silvestre, níspero, fruto del espino de mayo, pera, membrillo **Uvas y otra fruta pequeña trepadora de parra (exceptuando la fruta kiwi velludo):** Grosella espinosa, Fruta Kiwi Hardy, Uva Amur River, Pasionaria lila (Passiflora incarnata), Bayas Schisandra (Schisandra chinensis)

15

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

COMESTIBLES	
EN ESTAS PLANTAS (cont.)	**Verduras tuberosas y de bulbo:** Papa, camote, arracacha (Arracacia xanthorriza), sagú (Maranta arundinacea), alcachofa (china y de Jerusalén), achira (Canna edulis Ker-Gawler), yuca (Manihot esculenta) (amarga y dulce), chayote (Sechium edule) (raíz), chufa (Cyperus esculentus), sasheen, gengibre, lerén o topinambur (Calathea allouia), tanier (Caladium sagittæfolium), cúrcuma (Curcuma longa), jícama (Pachyrhizus erosus), batata (Dioscorea batatas) **Frutas con hueso:** Albaricoques, cereza (dulce y ácida), nectarinas, durazno, ciruela (Chickasaw, Damson, japonés), plumcot, ciruelas pasa (frescas) **Cucurbitáceas:** Chayote (fruta), calabaza china (melón chino para conserva), melón de agua (Citrullus lanatus), pepino, pepinillo, calabacín (comestible), mormordica spp., melón (híbrido y/o cultivares de Cucumis melo incluyendo melón Cucumis melo cantaloupensis, melón cantaloupensis, melón casaba, melón crenshaw, melón pershaw dorado, melón honeydew, melón honey balls, melón melon, melón pérsico, melón piña, melón Santa Claus, y melón culebra), calabaza, ayote (verano e invierno), sandía **Nueces de árboles:** Almendras, nuez de haya, nuez de Brasil, nuez de nogal, marañón, castaña, nuez de chinquapin, avellana, nuez de nogal americano, macadamia, pecana, pistacho, nuez (negra e inglesa) (persa) **Legumbres comestibles en vaina y arvejas suculentas y frijoles en vaina:** Frijol (*Phaseolus* spp.) (incluye frijol ayocote [Phaseolus coccineus], ejotes [Phaseolus vulgaris], frijol P. lunatus var. macrocarpus, o P. limensis, frijol Phaseolus lunatus [verde]); frijol (*Vigna* spp.) (incluye el frijol espárrago, ejote chino largo, guisantes polillá [Vigna aconitifolia], ejote de una yarda, frijol Vigna unguiculata, frijol salvaje [Vigna unguiculata], guisante del sur; frijol Canavalia ensiformis); haba (Vicia faba) (suculento); arveja (*Pisum* spp.) (incluye arveja enana, arveja de vaina comestible, arveja china o arveja de nieve [Pisum sativum], arveja “sugar snap”, arveja inglesa, arveja de jardín y arveja verde); soya (semilla inmadura); haba espada (Canavalia gladiata) y guandul (Cajanus cajan)

16

COMESTIBLES	
EN ESTAS PLANTAS (cont.)	**Fresas y otras bayas de baja estatura:** Gayuba (Arctostaphylos uva-ursi); arándano de arbusto bajo; mora de los pantanos (Rubus chamaemorus L.); arándano rojo; arándano de arbusto bajo o montañés (Vaccinium vitis-idaea); manzanas emúes (Kunzea pomifera F. Muell.); gaulteria, y cultivares, variedades y/o híbridos de éstas **Arándanos y otras bayas de arbusto y matorrales de tallo leñoso:** Baya de la Anoria (Aronia melanocarpa); arándano, de arbusto alto y bajo; grosellas; guayaba chilena; grosella, roja y negra; baya del saúco (Sambucus nigra); baya de agracejo (Berberis vulgaris); grosella espinosa; arándano, arbusto alto; madreselva comestible; zarzamora (Gaylussacia baccata); uvasella (Ribes nidigrolaria); baya de junio (Amelanchier alnifolia); arándano de arbusto bajo o montañés (Vaccinium vitis-idaea); grosella natie; baya del espino amarillo (Hippophae rhamnoides); y cultivares, variedades, y/o híbridos de éstos; y mora, frambuesa americana, frambuesa (negra y roja), frambuesa silvestre y cultivares, variedades y/o híbridos de éstas **Cebollas y otras verduras con bulbos:** Cebollines, hojas frescas; cebollines chinos, hojas frescas; bulbos del lirio daylily (género Hemerocallidaceae), elegans hosta (Hosta sieboldiana), hojas y bulbos de corona imperial (Fritillaria meleagris); ajo de bulbo grande, ajo de bulbo de serpiente; kurrat (Allium ampeloprasum var. kurrat); puerro (Allium cernuum); puerro; puerro silvestre, bulbo de lirio; cebolla Beltsville (Allium x proliferum [Moench] Schrad.), cebolla de bulbo, cebolla de bulbo chino; cebolla fresca, cebolla verde, cebolla de macrotallo, cebolla perla, cebolla bulbo de papa (Allium cepa L. var. aggregatum G. Don.); cebolla treetops (Allium x proliferum [Moench] Schrad. ex Willd.); cebolla Welsh tops (Allium fistulosum L.); bulbo de cebolla escalonia (Allium cepa var. aggregatum G. Don.), y hojas frescas y cultivares, variedades y/o híbridos de éstas
CUANDO APLICAR	Aplique tan pronto como detecte los problemas de insectos.

17

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

	Días de Espera Hasta la Cosecha	Días que Debe Esperar Antes de Otra Aplicación	Número Máximo de Aplicaciones por Temporada
Amaranto, rúcula silvestre, brócoli, col de Bruselas, repollo, cardo, coliflor, apio, celtuce (Lactuca sativa var. angustana), crisantemos (de hoja comestible, guirnalda), col verde, maíz salado, draba (jardín, meseta), diente de león, romaza, berenjena, endivia, hinojo de Florencia, uchuva, coles rizadas, colirrábano, lechuga (cabeza, hoja), mizuna, hojas verdes de mostaza, mostaza espinaca, plantas de sal, perejil, pepino, pimiento (morrón, chile, pimiento de cocina, pimiento, dulce), verdolaga (jardín, invierno), achicoria, nabicol (Brassica napus), ruibarbo, espinaca (hoja, de la vid, de Nueva Zelanda), acelga, tomatillo, tomate, papa, camote, arracacha (Arracacia xanthorrhiza), arracacha (Arracacia xanthorriza), sagú (Maranta arundinacea), alcachofa (china y de Jerusalén), achira (Canna edulis Ker-Gawler), yuca (Manihot esculenta) (amarga y dulce), chayote (Sechium edule) (raíz), chufa (Cyperus esculentus), sasheen, jengibre, lerén o topinambur (Calathea allouia), tanier (Caladium sagittæfolium), cúrcuma (Curcuma longa), jícama (Pachyrhizus erosus), batata (Dioscorea batatas), cebollines, hojas frescas; cebollines chinos, hojas frescas; bulbos del lirio daylily (género Hemerocallidaceae), cebolla Beltsville (Allium x proliferum,	7	7	4

	Días de Espera Hasta la Cosecha	Días que Debe Esperar Antes de Otra Aplicación	Número Máximo de Aplicaciones por Temporada
[Moench] Schrad.), cebolla de bulbo, cebolla de bulbo chino; cebolla fresca, cebolla verde, cebolla de macrotallo, cebolla perla, cebolla bulbo de papa (Allium cepa L. var. aggregatum G. Don.); cebolla treetops (Allium x proliferum [Moench] Schrad. ex Willd.); cebolla Welsh tops (Allium fistulosum L.); bulbo de cebolla escalonia (Allium cepa var. aggregatum G. Don.), y hojas frescas y cultivares, variedades y/o híbridos de éstas	7	7	4
Frutos de cítricos: Calamondín, híbridos de cítricos (chironja, tangelo, tangor), toronja, quinito, limón, lima, mandarina (tangerina), naranja (agria, dulce), pomelo, mandarina Satsuma	7	7	5
Frutos de pepitas: Manzana, manzano silvestre, níspero, fruto del espino de mayo, pera, membrillo	7	12	4
Uvas y otra fruta pequeña trepadora de parra (exceptuando la fruta kiwi velludo): Uva Amur River; grosella espinosa, fruta kiwi hardy, pasionaria lila (Passiflora incarnata), bayas Schisandra (Schisandra chinensis) y cultivares, variedades y/o híbridos de éstas	7	14	2
Frutas con hueso: Albaricoques, cereza (dulce y ácida), nectarinas, durazno, ciruela (Chickasaw, Damson, japonés), plumcot, ciruelas pasa (frescas)	7	10	4

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

	Días de Espera Hasta la Cosecha	Días que Debe Esperar Antes de Otra Aplicación	Número Máximo de Aplicaciones por Temporada
Cucurbitáceas: Chayote (fruta), calabaza china (melón chino para conserva), melón de agua (Citrullus lanatus), pepino, pepinillo, calabacín (comestible), mormordica spp., melón (híbrido y/o cultivares de Cucumis melo incluyendo melón Cucumis melo cantaloupensis, melón cantaloupensis, melón casaba, melón crenshaw, melón pershaw dorado, melón honeydew, melón honey balls, melón melon, melón pérsico, melón piña, melón Santa Claus, y melón culebra), calabaza, ayote (verano e invierno), sandía	0	5	5
Nueces de árboles: Almendras, nuez de haya, nuez de Brasil, nuez de nogal, marañón, castaña, nuez de chinquapin, avellana, nuez de nogal americano, macadamia, pecana, pistacho, nuez (negra e inglesa) (persa)	14	14	4
Legumbres comestibles en vaina y arvejas suculentas y frijoles en vaina: Frijol (*Phaseolus* spp.) (incluye frijol ayocote [Phaseolus coccineus], ejotes [Phaseolus vulgaris], frijol P. lunatus var. macrocarpus, o P. limensis, frijol Phaseolus lunatus [verde]); frijol (*Vigna* spp.) (incluye el frijol espárrago, ejote chino largo, guisantes polillá [Vigna aconitifolia], ejote de una yarda, frijol Vigna unguiculata, frijol salvaje [Vigna unguiculata], guisante del sur; frijol Canavalia ensiformis); haba (Vicia faba) (suculento); arveja (*Pisum* spp.) (incluye arveja enana, arveja de vaina	7	7	3

20

	Días de Espera Hasta la Cosecha	Días que Debe Esperar Antes de Otra Aplicación	Número Máximo de Aplicaciones por Temporada
comestible, arveja china o arveja de nieve [Pisum sativum], arveja “sugar snap”, arveja inglesa, arveja de jardín y arveja verde); soya (semilla inmadura); haba espada (Canavalia gladiata), y guandul (Cajanus cajan)	7	7	3
Fresas y otras bayas de baja estatura: Gayuba (Arctostaphylos uva-ursi); arándano de arbusto bajo; mora de los pantanos (Rubus chamaemorus L.); arándano rojo; arándano de arbusto bajo o montañés (Vaccinium vitis-idaea); manzanas emúes (Kunzea pomifera F. Muell.); gaulteria, y cultivares, variedades y/o híbridos de éstas	1	7	2
Arándanos y otras bayas de arbusto y matorral de tallo leñoso: Baya de la Anoria (Aronia melanocarpa); arándano, de arbusto alto y bajo; grosellas; guayaba chilena; grosella, roja y negra; baya del saúco (Sambucus nigra); baya de agracejo (Berberis vulgaris); grosella espinosa; arándano, arbusto alto; madreselva comestible; zarzamora (Gaylussacia baccata); uvasella (Ribes nidigrolaria); baya de junio (Amelanchier alnifolia); arándano de arbusto bajo o montañés (Vaccinium vitis-idaea); grosella natie; baya del espino amarillo (Hippophae rhamnoides); y cultivares, variedades, y/o híbridos de éstos; y mora, frambuesa americana, frambuesa (negra y roja), frambuesa silvestre y cultivares, variedades y/o híbridos de éstas	1	7	5

21

3.125”

3.125”

2.6875”

2.6875”

4.8125”

5.1875”

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO	Evite que el producto se congele. Mantenga el pesticida en su envase original. No lo coloque en envases de comidas o bebidas. Evite la contaminación de comestibles y alimentos para animales. Almacene el producto en un lugar fresco y seco, preferentemente en un área de almacenamiento bajo llave.
ELIMINACION	**Envase no rellenable. No vuelva a utilizar o llenar este envase.** **Si está vacío:** Colóquelo en la basura o recíclelo, si corresponde. **Si está parcialmente lleno:** Llame a su agencia de desechos sólidos local para obtener instrucciones sobre cómo eliminar el producto. Nunca elimine el producto no utilizado en drenajes internos o externos.

AVISOS DE PRECAUCION

PRECAUCION
RIESGOS PARA SERES HUMANOS (Y ANIMALES DOMESTICOS)

Es nocivo si se traga, inhala o se absorbe por la piel. Evite el contacto con los ojos, la piel o la ropa. Evite inhalar las emanaciones o vapores del rociado. Lávese bien con agua y jabón después de manipular el producto y antes de comer, beber, masticar chicle, usar tabaco o usar el baño. Debe quitarse y lavar la ropa contaminada antes de volver a usarla. El contacto prolongado o repetido con la piel puede causar reacciones alérgicas en algunos individuos.

PRIMEROS AUXILIOS

SI ENTRA EN CONTACTO CON LA PIEL	• Quítese la ropa contaminada. • Lave inmediatamente la piel con abundante agua durante 15 a 20 minutos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
EN CASO DE INHALAR	• Traslade a la persona a un lugar con aire fresco. • Si no respira, llame al 911 o a una ambulancia, luego realícele respiración artificial, de boca a boca, de ser posible. • Llame a un centro de control de envenenamientos o a un médico para obtener más recomendaciones sobre el tratamiento.
EN CASO DE INGERIR	• Consulte a un médico o a un centro de control de envenenamientos inmediatamente para obtener asesoramiento sobre el tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así se lo indique un médico o un centro de control de envenenamientos. • No administre nada por la boca a una persona que haya perdido el conocimiento.

Para emergencias médicas, llame las 24 horas del día al **1-800-225-2883**
Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a mano el envase o la etiqueta del producto.
También puede ponerse en contacto con el 1-800-225-2883 para obtener información médica de emergencia.

ADVERTENCIA PARA EL MEDICO: No existe un antídoto específico. Todos los tratamientos deben estar basados en los signos y síntomas de descompostura observados en el paciente. Puede haber ocurrido una exposición a otros materiales además de los de este producto.

3.125”

3.125”

BASE LABEL

2.75”

4.9375”

5.1875”

3.8125”

STORAGE AND DISPOSAL

STORAGE	Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place; preferably in a locked storage area.
DISPOSAL	**Non-refillable container. Do not reuse or refill this container.** **If empty**: Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

IF ON SKIN	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	• Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have the person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

For Medical Emergencies, Call 24 Hours a Day **1-800-225-2883**
Have the product container or label with you when calling a poison control center or doctor or going for treatment.
You may also contact 1-800-225-2883 for emergency medical treatment information.

NOTE TO PHYSICIAN: There is no specific antidote. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to other materials other than this product may have occurred.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This pesticide is toxic to wildlife. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treated area.

©2012 The ORTHO Group. World rights reserved.
Sold by **The ORTHO Group**
P.O. Box 190, Marysville, OH 43040
Product of Japan. LB2623

EPA Reg. No. 8033-107-239
EPA Est. 239-IA-3I, 239-MS-001M, 58996-MO-1A, 085652-OH-1W
Superscript is first letter of lot number.

PRESS TO RESEAL PRESIONE PARA VOLVER A CERRAR

EPA Reg No. 8033-107-239 (32 oz)
Ortho Flower, Fruit & Vegetable Insect Killer Ready-Spray II